

JOURNAL OF NUMBER THEORY 5, 203–214 (1973)

## On Character Sums and Class Numbers

MATTI JUTILA

*Department of Mathematics, University of Turku, Turku, Finland**Communicated by H. Halberstam*

Received March 27, 1971; revised August 21, 1971

A mean-value estimate for character sums with real characters is proved in an elementary way. The application is a proof of a conjecture on the error term in the asymptotic expression of the sum  $\sum_{d \leq X} h^k(-d)$ , where  $h(-d)$  is the class number of the quadratic field with the negative discriminant  $-d$ , and  $k$  is an integer  $\geq 2$ .

## INTRODUCTION

1. All real primitive characters are given by the Kronecker symbol  $\chi^{(d)}(n) = \left(\frac{d}{n}\right)$ , where  $d$  is a fundamental discriminant or 1, i.e., either  $d$  square-free and  $d \equiv 1 \pmod{4}$ , or  $d = 4N$ ,  $N$  square-free and  $N \equiv 2$  or  $3 \pmod{4}$ . The modulus of  $\chi^{(d)}$  is equal to  $|d|$ . Any real character is equivalent to a unique real primitive character and is, therefore, of the form  $\left(\frac{da^2}{n}\right)$ , where  $d$  is a fundamental discriminant or 1, and  $(a, d) = 1$ ,  $a$  square-free. Hence, all real nonprincipal characters are of the form  $\left(\frac{D}{n}\right)$ , with  $D$  belonging to a set  $\mathcal{D}$ , defined from the following conditions:

- (i)  $D$  is no square,
- (ii)  $D \equiv 1 \pmod{4}$ , or  $D = 4N$ ,  $N \equiv 1, 2$ , or  $3 \pmod{4}$ .

The object of this paper is to investigate sums of the shape

$$S(X, Y) = \sum'_{|D| \leq X} \left| \sum_{1 \leq n \leq Y} \left(\frac{D}{n}\right) \right|^2, \quad (1)$$

where, as also later,  $\sum'$  denotes a sum restricted to the set  $\mathcal{D}$ . By the classical Polya–Vinogradov estimate (shortly: P–V), the sum  $S(X, Y)$  is for  $X \geq 3$ ,  $Y \geq 1$  at most  $\ll X^2 \log^2 X$ . Our result is as follows:

THEOREM 1. For  $X \geq 3$ ,  $Y \geq 1$  we have uniformly

$$S(X, Y) \ll XY \log^8 X. \quad (2)$$

In the proof we need only classical arguments: a lemma of I. M. Vinogradov and its corollary (Lemmas 1 and 2), the law of reciprocity for real characters, and the estimate of Polya-Vinogradov.

2. It turns out that the sum over  $n$  in (1) may be restricted to fundamental discriminants without essential loss of sharpness of the estimate. Define

$$S_*^+(X, Y) = \sum'_{|D| \leq X} \left| \sum_{1 \leq n \leq Y}^* \left( \frac{D}{n} \right) \right|^2,$$

where  $\sum^*$  denotes a summation restricted to fundamental discriminants. Similarly, let  $S_*^-(X, Y)$  be defined as above but with  $n$  running over  $-Y \leq n \leq -1$ .

COROLLARY. For  $X \geq 3$ ,  $Y \geq 1$  we have uniformly

$$S_*^+(X, Y), S_*^-(X, Y) \ll XY \log^{10} X. \quad (3)$$

3. As an application of the corollary, we shall prove a mean-value theorem for  $h^k(-d)$ , with  $h(-d)$  the class number of the quadratic field  $\mathbb{Q}(\sqrt{-d})$  with the negative discriminant  $-d$  and  $k \geq 1$  a fixed integer.

THEOREM 2. Let  $k \geq 1$  be an integer. Then for  $X \geq 3$  we have

$$\sum_{1 \leq d \leq X}^* h^k(-d) = a(k) X^{\frac{1}{2}(k+2)} (1 + O(X^{-1/2} (\log X)^{c_1(k)})),$$

where  $a(k)$ ,  $c_1(k)$  and the  $O$ -constant depend on  $k$  only.

Using the Dirichlet formula  $h(-d) = \pi^{-1} |d|^{1/2} L(1, \chi^{(-d)})$ , valid for  $d > 4$  (see [3], Chap. 5, Sec. 4), Theorem 2 is by partial summation obtained from the following

THEOREM 3. With  $k$  and  $X$  as in Theorem 2, we have

$$\sum_{d \in [e, eX]}^* L^k(1, \chi^{(d)}) = b(k)X + O(X^{1/2} (\log X)^{c_2(k)}),$$

where  $e = \pm 1$ , and  $b(k)$ ,  $c_2(k)$  and the  $O$ -constant depend on  $k$  only.

## NOTES

4. Wolke has in [9] estimated a double sum for quadratic characters, using the "large-sieve" inequality of Linnik–Bombieri–Gallagher. It may be noted that, as an easy consequence of Theorem 1, Lemma 1 of [9] can be improved in the case  $y \geq x^{1/2+\epsilon}$ .

5. Sums closely related to the one considered in Theorem 2 have been investigated in the case  $k = 1$  by I. M. Vinogradov [6], [7] and in the case  $k \geq 2$  by Lavrik [4]; Barban [1], [2]; Wolke [9]; and also by some other writers. These papers concern the  $k$ 'th power of the class number of the properly primitive positive binary quadratic forms with the negative determinant  $-d$ . In other words, the forms in question are of the type  $ax^2 + 2bxy + cy^2$  with  $a > 0$ ,  $(a, 2b, c) = 1$ ,  $-d = b^2 - ac < 0$ . An estimate of the same strength as Theorem 2 can be proved for this class number sum.

## A LEMMA OF I. M. VINOGRADOV

6. The following well-known lemma is proved in [8], pp. 32–34.

LEMMA 1. *Let  $\alpha$ ,  $\beta$ , and  $\Delta$  be real numbers, satisfying*

$$0 < \Delta < \frac{1}{2}, \quad \Delta \leq \beta - \alpha \leq 1 - \Delta.$$

*Then there exists a periodic function  $\psi(x)$  with period 1, satisfying*

- (i)  $\psi(x) = 1$  in the interval  $\alpha + \frac{1}{2}\Delta \leq x \leq \beta - \frac{1}{2}\Delta$ ,
- (ii)  $\psi(x) = 0$  in the interval  $\beta + \frac{1}{2}\Delta \leq x \leq 1 + \alpha - \frac{1}{2}\Delta$ ,
- (iii)  $0 \leq \psi(x) \leq 1$  in the intervals  $\alpha - \frac{1}{2}\Delta \leq x \leq \alpha + \frac{1}{2}\Delta$ ,  $\beta - \frac{1}{2}\Delta \leq x \leq \beta + \frac{1}{2}\Delta$ ,
- (iv)  $\psi(x)$  has the Fourier expansion

$$\psi(x) = \beta - \alpha + \sum_{m=1}^{\infty} (a_m e^{2\pi i m x} + \bar{a}_m e^{-2\pi i m x}), \quad (4)$$

where

$$a_m = (2m\pi i)^{-1} (e^{-2\pi i m \alpha} - e^{-2\pi i m \beta}) \left\{ \frac{\sin(\frac{1}{2}\pi m \Delta)}{\frac{1}{2}\pi m \Delta} \right\}^2, \quad (5)$$

$$|a_m| \leq \min\{\beta - \alpha, m^{-1}, \Delta^{-2} m^{-3}\}, \quad (6)$$

(v) if  $\alpha$  and  $\Delta$  are fixed but  $\beta$  variable in the interval  $B = [\alpha + \Delta, \alpha + 1 - \Delta]$ , and the function  $\psi(x)$ , corresponding to a particular  $\beta$ , is denoted by  $\psi_\beta(x)$ , then we have

$$\psi_{\beta'}(x) = \psi_\beta(x + \beta - \beta') \quad (7)$$

for all  $\beta, \beta' \in B$ , and for  $\beta' - \min(\beta - \alpha - \frac{1}{2}\Delta, \beta' - \alpha - \frac{1}{2}\Delta) \leq x \leq \beta' + \min(1 + \alpha - \frac{1}{2}\Delta - \beta, 1 + \alpha - \frac{1}{2}\Delta - \beta')$ .

Indeed, take  $r = 2$  in Lemma 2 of [8]. The additional assertion (v) follows from the method of construction of the function  $\psi(x)$ .

For future reference, it is convenient to state some further properties of  $\psi(x)$ .

LEMMA 2. Denote by  $\psi(x, u)$  the function  $\psi(x)$  of Lemma 1 with the following choice of the parameters:

$$\Delta = X^{-1/2}, \quad \alpha = \frac{1}{2}\Delta, \quad \beta = Yu^{-1} + \frac{1}{2}\Delta = \beta(u), \quad (8)$$

where

$$X^{1/2} \leq Y \leq \frac{1}{4}X, \quad \frac{1}{2}X \leq u \leq X, \quad X > 100. \quad (9)$$

Then

$$\frac{\partial \psi(x, u)}{\partial x} \ll \Delta^{-1}. \quad (10)$$

Further,

$$\frac{d\psi\left(\frac{n}{u}, u\right)}{du} \ll X^{-1} \quad (11)$$

if

$$1 \leq n \leq X^{1/2}, \quad \text{or} \quad Y \leq n \leq Y + X^{1/2}. \quad (12)$$

Finally, if  $a_{m,u}$  are the Fourier coefficients of the expansion of the function  $\psi(x, u)$ , then

$$\frac{da_{m,u}}{du} \ll \min\{YX^{-2}, YX^{-1}m^{-2}\}. \quad (13)$$

The estimates (10), (11), and (13) hold uniformly with respect to all parameters and variables.

*Proof.* For the proof of (10), note that the expansion (4) can be differentiated termwise, and use (6).

The function  $\psi(x, u)$  is, for  $0 \leq x \leq 2\Delta$ , independent of  $u$ ; hence for  $1 \leq n \leq X^{1/2}$  we have by (10), (8), and (9) the estimate (11).

For the remaining case  $Y \leq n \leq Y + X^{1/2}$  of (11), we use (v) of Lemma 1 with  $\beta = (Y/u) + \frac{1}{2}\Delta$ ,  $\beta' = [Y/(u + \eta)] + \frac{1}{2}\Delta$ . We have

$$\begin{aligned} \frac{d\psi\left(\frac{n}{u}, u\right)}{du} &= \lim_{\eta \rightarrow 0} \eta^{-1} \left( \psi\left(\frac{n}{u + \eta}, u + \eta\right) - \psi\left(\frac{n}{u}, u\right) \right) \\ &= \lim_{\eta \rightarrow 0} \eta^{-1} \left( \psi\left(\frac{n}{u + \eta} + \frac{Y}{u} - \frac{Y}{u + \eta}, u\right) - \psi\left(\frac{n}{u}, u\right) \right) \\ &\ll \lim_{\eta \rightarrow 0} \eta^{-1} \Delta^{-1}(n - Y) \left( \frac{1}{u + \eta} - \frac{1}{u} \right) \ll X^{-1}, \end{aligned}$$

and (11) is established.

The estimate (13) is obtained from (5) by a simple calculation.

#### PROOF OF THEOREM 1

7. The case  $Y \geq \frac{1}{4}X$  is clear by P-V, and the case  $Y \leq X^{1/2}$  is also easy, as seen from the following calculation. First,

$$S(X, Y) \leq \sum_{\substack{r, s \leq Y \\ rs \neq a^2}} \sum'_{|D| \leq X} \left( \frac{D}{rs} \right) + \sum_{n^2 \leq Y^2} \tau(n^2) \sum'_{|D| \leq X} 1. \quad (14)$$

To estimate the second double sum, note that  $\tau(n^2) \leq \tau^2(n)$ , and that

$$\sum_{n \leq Y} \tau^2(n) \ll Y \log^3 Y,$$

whence the resulting contribution to  $S(X, Y)$  is  $\ll XY \log^3 Y$ .

Modify the first double sum removing the condition (i) from the definition of the set  $\mathcal{D}$ . This gives rise to an error  $\ll Y^2 X^{1/2} \leq XY$  which can be neglected.

In the resulting double sum, let first  $rs$  be fixed and consider the values of the symbol

$$\left( \frac{D}{rs} \right).$$

For the values of  $D$  in question, we have

$$\left( \frac{D}{rs} \right) = \chi(D),$$

where  $\chi$  is a nonprincipal character to a modulus  $\leq 4rs$ , having a con-

ductor different from 4. Hence the  $D$ -sum can be expressed by a finite number of character sums with nonprincipal characters to a modulus  $\leq 16rs$ . So, P-V, the double sum in consideration is

$$\ll \sum_{r,s \leq Y} (rs)^{1/2} \log Y \ll Y^3 \log X \leq XY \log X.$$

Hence Theorem 1 is found to hold in the case  $Y \leq X^{1/2}$ .

8. Now we proceed to the more difficult case  $X^{1/2} \leq Y \leq \frac{1}{4}X$ . Since we shall apply Lemma 2 with  $u = |D|$ , it is convenient to restrict  $|D|$  to the interval  $[\frac{1}{2}X, X]$ , and the sum over this interval being estimated, apply the result with  $\frac{1}{2}X$  in place of  $X$ , etc. Furthermore, we first restrict  $D$  to the sequence of the fundamental discriminants.

The first step is to express the inner sum (e.g.,  $S^{(D)}$ ) of  $S(X, Y)$  in terms of the function  $\psi(x, u)$  of Lemma 2 as follows:

$$\begin{aligned} S^{(D)} &= \sum_{1 \leq n \leq Y} \left( \frac{D}{n} \right) = \sum_{1 \leq n \leq |D|} \psi \left( \frac{n}{|D|}, |D| \right) \left( \frac{D}{n} \right) \\ &\quad + \sum_{1 \leq n \leq X^{1/2}} \left( 1 - \psi \left( \frac{n}{|D|}, |D| \right) \right) \left( \frac{D}{n} \right) \\ &\quad - \sum_{Y \leq n \leq Y+X^{1/2}} \psi \left( \frac{n}{|D|}, |D| \right) \left( \frac{D}{n} \right) \\ &= S_1^{(D)} + S_2^{(D)} + S_3^{(D)}. \end{aligned} \quad (15)$$

Hence, writing  $S$  for the part of  $S(X, Y)$  in consideration, we have

$$S = \sum_{\frac{1}{2}X \leq |D| \leq X}^* |S^{(D)}|^2 \ll \sum_{i=1}^3 \sum_{\frac{1}{2}X \leq |D| \leq X}^* |S_i^{(D)}|^2 = S_1 + S_2 + S_3. \quad (16)$$

9. *Treatment of the sum  $S_1$ .* Using the representation

$$\psi(x, u) = \beta(u) - \alpha + \sum_{m=1}^{\infty} (a_{m,u} e^{2\pi i m x} + \bar{a}_{m,u} e^{-2\pi i m x})$$

and the primitivity of the character  $\left( \frac{D}{n} \right)$  in (15), we obtain

$$\begin{aligned} S_1^{(D)} &= \tau_D \sum_{m=1}^{\infty} \left( a_{m,|D|} \left( \frac{D}{m} \right) + \bar{a}_{m,|D|} \left( \frac{D}{-m} \right) \right) \\ &= \tau_D \left( \sum_{m \leq X^{1/2}} + \sum_{m > X^{1/2}} \right) = S_{11}^{(D)} + S_{12}^{(D)}, \end{aligned}$$

where  $\tau_D$  is a Gaussian sum. So

$$S_1 \ll \sum_{\frac{1}{2}X \leq |D| \leq X}^* (|S_{11}^{(D)}|^2 + |S_{12}^{(D)}|^2) = S_{11} + S_{12}.$$

Since  $|\tau_D|^2 = |D|$ , we have further, allowing  $D$  now run over  $\mathcal{D}$ ,

$$\begin{aligned} S_{11} &\ll \sum'_{\frac{1}{2}X \leq |D| \leq X} |D| \left| \sum_{m \leq X^{1/2}} a_{m,|D|} \left( \frac{D}{m} \right) \right|^2 \\ &= \sum_{r, s \leq X^{1/2}} \sum'_{\frac{1}{2}X \leq |D| \leq X} |D| a_{r,|D|} \bar{a}_{s,|D|} \left( \frac{D}{rs} \right), \end{aligned} \quad (17)$$

and analogously for  $S_{12}$ .

To estimate the sum  $S_{11}$ , we consider separately two cases according to whether  $rs$  is a square or not.

(A) *rs a square.* By (6) and (8), the contribution of such pairs  $(r, s)$  to the sum on the right of (17) is

$$\ll X^2 \sum_{r, s \leq X^{1/2}} q(rs) \min(YX^{-1}, r^{-1}) \min(YX^{-1}, s^{-1}), \quad (18)$$

with  $q(n) = 1$  if  $n$  is a square, and  $q(n) = 0$  otherwise.

Split up the double sum (18) into  $\ll \log^2 X$  parts of the type

$$\sum(U, V) = \sum_{\substack{U \leq r \leq U' \\ V \leq s \leq V'}},$$

with  $U' \leq 2U$ ,  $V' \leq 2V$ . Now by (18) obviously

$$\sum(U, V) \ll X^2 \sum_{UV \leq n^2 \leq 4UV} \tau(n^2) \min(YX^{-1}, U^{-1}) \min(YX^{-1}, V^{-1}). \quad (19)$$

Considering separately the four cases combined from  $U \leq XY^{-1}$ ,  $U > XY^{-1}$ , and  $V \leq XY^{-1}$ ,  $V > XY^{-1}$ , it is easily seen that in any case

$$\sum(U, V) \ll XY \log^3 X,$$

and so all sums of this kind contribute to  $S_{11}$  together  $\ll XY \log^5 X$ .

(B) *rs not a square.* In this case we estimate the inner sums in (17) by partial summation. To this end, we introduce the functions

$$f_{r,s}(u) = ua_{r,u} \bar{a}_{s,u}, \quad (20)$$

$$g_{r,s}(u) = \sum'_{\frac{1}{2}X \leq |D| \leq u} \left( \frac{D}{rs} \right). \quad (21)$$

Let  $p(n) = 0$  if  $n$  is a square, and  $p(n) = 1$  otherwise. Then the part in consideration of the double sum in (17) is by a well-known formula equal to

$$\sum_{r,s \leq X^{1/2}} p(rs) \left\{ g_{r,s}(X) f_{r,s}(X) - \int_{\frac{1}{2}X}^X g_{r,s}(u) f'_{r,s}(u) du \right\}. \quad (22)$$

Now note that by (6)

$$f_{r,s}(X) \ll X r^{-1} s^{-1}, \quad (23)$$

by P-V (as in Sec. 7)

$$g_{r,s}(u) \ll (rs)^{1/2} \log X + X^{1/2} \ll X^{1/2} \log X, \quad (24)$$

and finally by (13) and (6)

$$f'_{r,s}(u) \ll YX^{-1}(|a_{r,u}| + |a_{s,u}|) \ll YX^{-1}(r^{-1} + s^{-1}). \quad (25)$$

Using (23)–(25), we obtain for (22) the estimate

$$\ll \sum_{r,s \leq X^{1/2}} X^{1/2} \log X (X r^{-1} s^{-1} + Y(r^{-1} + s^{-1})) \ll XY \log^3 X.$$

We have proved that  $S_{11} \ll XY \log^5 X$ . The above method works for the sum  $S_{12}$  as well. Now  $r$  and  $s$  run over  $r > X^{1/2}, s > X^{1/2}$ . The Case (A) gives a contribution

$$\ll X^2 \sum_{n^2 > X} \tau(n^2) X^2 n^{-6} \ll X^{3/2} \log^3 X \ll XY \log^3 X.$$

For the Case (B) note that in place of (23)–(25) we now have the estimates  $f_{r,s}(X) \ll X^3 r^{-3} s^{-3}$ ,  $g_{r,s}(u) \ll (rs)^{1/2} \log(rs)$ ,  $f'_{r,s}(u) \ll XY r^{-2} s^{-2} (r^{-1} + s^{-1})$ , and by partial summation we obtain a contribution  $\ll XY \log X$ .

To summarize,  $S_{12} \ll XY \log^3 X$ , hence by the above estimate for  $S_{11}$  we have finally

$$S_1 \ll XY \log^5 X. \quad (26)$$

**10. Treatment of the sums  $S_2$  and  $S_3$ .** We shall consider in detail the sum  $S_3$  only since the case of  $S_2$  is quite analogous.

By (15) and (16) we have

$$S_3 \leq \sum_{Y \leq r, s \leq Y+X^{1/2}} \sum'_{\frac{1}{2}X \leq |D| \leq X} \psi\left(\frac{r}{|D|}, |D|\right) \psi\left(\frac{s}{|D|}, |D|\right) \left(\frac{D}{rs}\right).$$

Again distinguish two cases according to whether  $rs$  is a square or not. The squares give a contribution  $\ll X^{3/2} \log^3 X \ll XY \log^3 X$ , as easily calculated.



The remaining sum is by partial summation equal to

$$\sum_{Y \leq r, s \leq Y+X^{1/2}} p(rs) \left\{ \psi\left(\frac{r}{X}, X\right) \psi\left(\frac{s}{X}, X\right) g_{r,s}(X) - \int_{\frac{1}{2}X}^X g_{r,s}(u) \frac{d}{du} \left( \psi\left(\frac{r}{u}, u\right) \psi\left(\frac{s}{u}, u\right) \right) du \right\}.$$

By (11) and P-V, this is

$$\ll \sum_{Y \leq r, s \leq Y+X^{1/2}} (rs)^{1/2} \log X \ll XY \log X.$$

Hence,  $S_3 \ll XY \log^3 X$ , and similarly for  $S_2$ . So, in view of (26) and (16), we have  $S \ll XY \log^5 X$ , and hence

$$\sum_{|D| \leq X}^* \left| \sum_{1 \leq n \leq Y} \left( \frac{D}{n} \right) \right|^2 \ll XY \log^6 X. \quad (27)$$

**11. Completion of the proof.** In view of (27), we only have to estimate the contribution of the nonfundamental values of  $D$  to the sum  $S(X, Y)$ .

In general, if  $\left(\frac{d}{n}\right)$  is the primitive character, equivalent to  $\left(\frac{D}{n}\right)$ , and  $D = dd_1$ , we have

$$\begin{aligned} \left| \sum_{1 \leq n \leq Y} \left( \frac{D}{n} \right) \right|^2 &= \left| \sum_{\substack{1 \leq n \leq Y \\ (n, d_1)=1}} \left( \frac{d}{n} \right) \right|^2 = \left| \sum_{\delta/d_1} \mu(\delta) \left( \frac{d}{\delta} \right) \sum_{1 \leq n \leq Y\delta^{-1}} \left( \frac{d}{n} \right) \right|^2 \\ &\leq \tau(d_1) \sum_{\delta/d_1} \left| \sum_{1 \leq n \leq Y\delta^{-1}} \left( \frac{d}{n} \right) \right|^2. \end{aligned}$$

Hence for fixed values of  $d$  and  $\delta$  with  $|d| \delta \leq X$  we have to  $S(X, Y)$  a contribution

$$\leq \left| \sum_{1 \leq n \leq Y\delta^{-1}} \left( \frac{d}{n} \right) \right|^2 \sum_{\substack{d_1 \leq X|d|^{-1} \\ \delta/d_1}} \tau(d_1) \ll \left| \sum_{1 \leq n \leq Y\delta^{-1}} \left( \frac{d}{n} \right) \right|^2 \tau(\delta) X |d|^{-1} \delta^{-1} \log X.$$

Now a summation over  $|d| \leq X\delta^{-1}$  for a fixed  $\delta$  gives by (27) a contribution  $\ll XY\delta^{-2}\tau(\delta) \log^8 X$ . Finally, a summation over  $\delta$  accomplishes the proof of Theorem 1.

## PROOF OF THE COROLLARY

12. Consider  $S_*^+(X, Y)$ . Let generally in a sum  $\sum_n''$  the number  $n$  run over the integers  $n = \nu a^2$ , where  $a$  is odd and  $\nu$  a fundamental discriminant. Then obviously

$$\begin{aligned} \left| \sum_{1 \leq n \leq Y}^* \left( \frac{D}{n} \right) \right| &= \left| \sum_{1 \leq n \leq Y}'' \left( \sum_{\substack{\delta^2 | n \\ 2 \nmid \delta}} \mu(\delta) \right) \left( \frac{D}{n} \right) \right| \leq \sum_{\substack{\delta \leq Y^{1/2} \\ 2 \nmid \delta}} \left| \sum_{1 \leq n \leq Y}'' \left( \frac{D}{n} \right) \right| \\ &\leq \sum_{\substack{\delta \leq Y^{1/2} \\ 2 \nmid \delta}} (\Sigma_1^{(D, \delta)} + \Sigma_2^{(D, \delta)}), \end{aligned} \quad (28)$$

where

$$\Sigma_1^{(D, \delta)} = \left| \sum_{\substack{1 \leq n \leq Y \\ 4\delta^2 | n}}'' \left( \frac{D}{n} \right) \right|,$$

and  $\Sigma_2^{(D, \delta)}$  a similar sum with the conditions  $n \equiv 1 \pmod{4}$ ,  $\delta^2 | n$  instead of the condition  $4\delta^2 | n$ .

In  $\Sigma_1^{(D, \delta)}$ ,  $n$  runs over the numbers  $n = 4\delta^2 k$ ,  $k \equiv 2, 3 \pmod{4}$  and in  $\Sigma_2^{(D, \delta)}$  over the numbers  $n = \delta^2 k$ ,  $k \equiv 1 \pmod{4}$ . Hence it is easily seen that

$$\begin{aligned} \Sigma_1^{(D, \delta)} &\leq \left| \sum_{k \leq Y/4\delta^2} \left( \frac{D}{k} \right) \right| + \left| \sum_{k \leq Y/16\delta^2} \left( \frac{D}{k} \right) \right| \\ &\quad + \left| \sum_{k \leq Y/4\delta^2} \chi_4(k)(1 + \chi_4(k)) \left( \frac{D}{k} \right) \right|, \end{aligned} \quad (29)$$

$$\Sigma_2^{(D, \delta)} \leq \left| \sum_{k \leq Y/\delta^2} \chi_4(k)(1 + \chi_4(k)) \left( \frac{D}{k} \right) \right|, \quad (30)$$

where  $\chi_4$  is the primitive character  $\pmod{4}$ .

From (28)–(30), we see that the proof of the estimate of  $S_*^+(X, Y)$  is reduced to the proof of the estimate

$$\sum_{|D| \leq X} \left( \sum_{\substack{\delta \leq Y^{1/2} \\ 2 \nmid \delta}} \left| \sum_{n \leq Y\delta^{-2}} \left( \frac{D}{n} \right) \right| \right)^2 \ll XY \log^{10} X.$$

But this is easily verified by Schwarz's inequality and Theorem 1. Since the case of  $S_*^-(X, Y)$  is quite analogous, the Corollary is verified.

## PROOF OF THEOREM 3

## 13. Starting from the formula

$$L(1, \chi^{(d)}) = \sum_{n \leq X} \frac{\left(\frac{d}{n}\right)}{n} + O(X^{-1/2} \log X),$$

valid for all  $d$  with  $|d| \leq X$ , we have

$$L^k(1, \chi^{(d)}) = \sum_{n \leq X^k} \frac{\tau_k'(n) \left(\frac{d}{n}\right)}{n} + O_k(X^{-1/2} \log^k X),$$

where  $\tau_k'(n) = \tau_k(n)$  for  $n \leq X$ , and  $\tau_k'(n) \leq \tau_k(n)$  for  $n > X$ . So, on summation over  $d$ , we obtain

$$\begin{aligned} \sum_{d \in [e, eX]}^* L^k(1, \chi^{(d)}) &= \sum_{n^2 \leq X} \frac{\tau_k(n^2)}{n^2} \sum_{\substack{d \in [e, eX] \\ (d, n)=1}}^* 1 + O\left(\sum_{n^2 > X} \frac{\tau_k(n^2)}{n^2} X\right) \\ &\quad + O_k(X^{1/2} \log^k X) + O\left(\sum_{\substack{n \leq X^k \\ n \neq a^2}} \frac{\tau_k(n) |S(n)|}{n}\right), \end{aligned} \quad (31)$$

where

$$S(n) = \sum_{d \in [e, eX]}^* \left(\frac{d}{n}\right).$$

Consider the main term first. It is easily verified that

$$\sum_{\substack{d \in [e, eX] \\ (d, n)=1}}^* 1 = A(n)X + O(X^{1/2} \tau(n))$$

on expressing the sum by the Mobius function. Hence the main term in (31) is

$$b(k)X + O_k(X^{1/2}),$$

where

$$b(k) = \sum_{n=1}^{\infty} A(n) \tau_k(n^2) n^{-2}.$$

To prove Theorem 3, we have to prove an estimate for the last term in (31), and the result

$$\sum_{\substack{n \leq X^k \\ n \neq a^2}} \frac{|S(n)|}{n} = O_k(X^{1/2} \log^6 X) \quad (32)$$

would suffice. But by the reciprocity law for real characters, we can write the sum  $S(n)$  as a sum of the type considered in  $S_*^+(X, Y)$  and  $S_*^-(X, Y)$ . Hence, by Schwarz's inequality and the Corollary, the estimates (32) is established, and the proof of Theorem 3 is complete.

## REFERENCES

1. M. B. BARBAN, "Large sieve" of Yu. V. Linnik and a limit theorem for the class number of an imaginary quadratic field, *Izv. Akad. Nauk SSSR, Ser. Mat.* **26** (1962), 573-580 (in Russian).
2. M. B. BARBAN, On a density hypothesis of E. Bombieri, *Dokl. Akad. Nauk SSSR* **172** (1967), 999-1000 (in Russian).
3. Z. I. BOREVICH AND I. R. SHAFAREVICH, "Number Theory," Academic Press, New York and London, 1966.
4. A. F. LAVRIK, On the problem of the distribution of values of the class number of properly primitive quadratic forms of negative determinant, *Izv. Akad. Nauk UzSSR, Ser. Fiz.-Mat.* (1959), No. 1, 81-90 (in Russian).
5. A. F. LAVRIK, Double sums with quadratic characters, *Dokl. Akad. Nauk SSSR* **186** (1969), 19-21 (in Russian).
6. I. M. VINOGRADOV, A new method for obtaining asymptotic expressions for arithmetical functions, *Izv. Ross. Akad. Nauk* (1917), 1347-1378 (in Russian).
7. I. M. VINOGRADOV, On the problem of the number of integral points in a given region, *Izv. Akad. Nauk SSSR, Ser. Mat.* **24** (1960), 777-786 (in Russian).
8. I. M. VINOGRADOV, "The method of trigonometrical sums in the theory of numbers," Interscience Publishers, London-New York, 1955.
9. DIETER WOLKE, Moments of the Number of Classes of Primitive Quadratic Forms with Negative Discriminant, *J. Number Theory* **1** (1969), 502-511.